

FarmLens Ltd

Website: farmlens.africa | App: app.farmlens.africa | Headquarters: Nairobi, Kenya



Crop details

Oregano

Origanum vulgare

Family: Lamiaceae

Categories

Vegetables

Spices & Condiments

Generated: 2026-04-11 10:16

Quick stats

Family	Lamiaceae
Typical harvest	6.5 t/ha
Varieties	1
Pests and diseases	2
Seasons	1

Crop profile

Growth habit	perennial
Days to harvest	90
Main uses	Leaves for seasoning, drying, and herb blends.
Pollination	insect
Origin and where it grows	Adapted to dry highland herb production and kitchen gardens.

Weather, soil and spacing

Best temperature	14 - 24 °C
Rainfall	450 - 800 mm/yr
Altitude	800 - 2800 m
Best pH	6 - 7
Soil type	Fertile well-drained loam rich in organic matter.
Row spacing	35 cm
Plant spacing	20 cm
Planting depth	1.5 cm
Seed rate	3 kg/ha

Simple notes for farmers

About the crop: This crop is perennial; once planted it can keep producing for many years. Harvest typically starts about 90 days after planting.

Main use: Farmers mostly grow this crop for leaves for seasoning, drying, and herb blends..

Pollination: Mainly insect; healthy flowers and pollinators improve fruit set.

Where it grows: Adapted to dry highland herb production and kitchen gardens.. Grouped under: Vegetables, Spices & Condiments.

Best climate: 14 - 24 °C; 450 - 800 mm/yr; up to about 2800 m a.s.l.

Soil: Best at pH 6 - 7; fertile well-drained loam rich in organic matter..

Farmer guide (Mwongozo wa Mkulima)

<u>Planting</u>	Establish Oregano in a fine weed-free seedbed and keep emergence moisture steady.
<u>Transplanting</u>	Direct seed or transplant depending on production system.
<u>Irrigation</u>	Maintain even soil moisture for steady Oregano growth and quality.
<u>Fertigation</u>	Use split nitrogen and potassium for market-quality Oregano.
<u>Pest scouting</u>	Scout Oregano weekly for chewing pests, sap suckers, and foliar diseases.
<u>Pruning and training</u>	No pruning required unless sanitation or staking is needed.
<u>Harvest</u>	Harvest Oregano in the cool hours and handle gently to preserve aroma.
<u>Postharvest</u>	Cool and shade Oregano promptly after harvest.

Nutrient schedule (Mbolea kwa Hatua)

#	Stage	DAP	Product	Rate	Targets (kg/ha)	Notes
1	Basal	0	NPK 17-17-17	200 kg/ha	N: 34, P ₂ O ₅ : 34, K ₂ O: 34	Basal fertilizer for Oregano.
2	Topdress	21	CAN	100 kg/ha	N: 26, P ₂ O ₅ : N/A, K ₂ O: N/A	Support active Oregano vegetative growth.

Nutrient requirements

Nutrient	Stage	Amount	Unit
N	Basal	35	kg/ha
P ₂ O ₅	Basal	30	kg/ha
K ₂ O	Basal	35	kg/ha
N	Topdress	25	kg/ha
K ₂ O	Topdress	20	kg/ha

Field images



Varieties

Name	Country	Maturity (days)	Traits
Greek Oregano	KE	90	High aroma and good drying quality.

Fertilizer recommendations

<u>Stage</u>	<u>Product</u>	<u>Rate</u>	<u>Notes</u>
Planting	Well-rotted manure	5000	Improve soil structure before Oregano planting.
Vegetative growth	CAN	100	Split topdress for Oregano production.

Pests and diseases

<u>Name</u>	<u>Type</u>	<u>Symptoms</u>	<u>Management</u>
Aphids	pest	Crowding on tender shoots and reduced growth.	Use clean fields and selective control if needed.
Powdery mildew	disease	White powdery growth on leaves.	Improve airflow and avoid overcrowding.

Yields

<u>System</u>	<u>Typical</u>	<u>Min</u>	<u>Max</u>	<u>Notes</u>
Managed fresh-market production	6.5	4.6	9.8	Typical marketable Oregano yield under irrigated or well-managed conditions.

Season calendars

<u>Country</u>	<u>Region</u>	<u>Planting</u>	<u>Harvest</u>
KE	Highland Vegetable Zones	Mar-Apr or Oct-Nov	Year-round depending on irrigation

Region suitability

<u>Country</u>	<u>Region</u>	<u>Suitability</u>
KE	Highland Vegetable Zones	High

Source: **FarmLens Ltd** - farmlens.africa and app.farmlens.africa. Headquarters: Nairobi, Kenya. This guide was generated from the FarmLens database.