

FarmLens Ltd

Website: farmlens.africa | App: app.farmlens.africa | Headquarters: Nairobi, Kenya



Crop details

Parsnip

Pastinaca sativa

Family: Apiaceae

Categories

Vegetables

Generated: 2026-04-11 10:16

Quick stats

Family	Apiaceae
Typical harvest	16.0 t/ha
Varieties	1
Pests and diseases	2
Seasons	1

Crop profile

Growth habit	annual
Days to harvest	120
Main uses	Sweetened storage root for soups, roasting, and fresh markets.
Pollination	insect
Origin and where it grows	Best in cool highland areas with deep friable soils.

Weather, soil and spacing

Best temperature	10 - 20 °C
Rainfall	450 - 800 mm/yr
Altitude	800 - 2800 m
Best pH	6 - 7
Soil type	Fertile well-drained loam rich in organic matter.
Row spacing	30 cm
Plant spacing	10 cm
Planting depth	1.5 cm
Seed rate	5 kg/ha

Simple notes for farmers

About the crop: This crop is annual; it grows and is harvested in one season. Harvest typically starts about 120 days after planting.

Main use: Farmers mostly grow this crop for sweetened storage root for soups, roasting, and fresh markets..

Pollination: Mainly insect; healthy flowers and pollinators improve fruit set.

Where it grows: Best in cool highland areas with deep friable soils.. Grouped under: Vegetables.

Best climate: 10 - 20 °C; 450 - 800 mm/yr; up to about 2800 m a.s.l.

Soil: Best at pH 6 - 7; fertile well-drained loam rich in organic matter..

Farmer guide (Mwongozo wa Mkulima)

<u>Planting</u>	Establish Parsnip in a fine weed-free seedbed and keep emergence moisture steady.
<u>Transplanting</u>	Direct seed or transplant depending on production system.
<u>Irrigation</u>	Maintain even soil moisture for steady Parsnip growth and quality.
<u>Fertigation</u>	Use split nitrogen and potassium for market-quality Parsnip.
<u>Pest scouting</u>	Scout Parsnip weekly for chewing pests, sap suckers, and foliar diseases.
<u>Pruning and training</u>	No pruning required unless sanitation or staking is needed.
<u>Harvest</u>	Harvest Parsnip when roots size up well before they become fibrous or pithy.
<u>Postharvest</u>	Cool and shade Parsnip promptly after harvest.


Nutrient schedule (Mbolea kwa Hatua)

<u>#</u>	<u>Stage</u>	<u>DAP</u>	<u>Product</u>	<u>Rate</u>	<u>Targets (kg/ha)</u>	<u>Notes</u>
1	Basal	0	NPK 17-17-17	200 kg/ha	N: 34, P ₂ O ₅ : 34, K ₂ O: 34	Basal fertilizer for Parsnip.
2	Topdress	21	CAN	100 kg/ha	N: 26, P ₂ O ₅ : N/A, K ₂ O: N/A	Support active Parsnip vegetative growth.

Nutrient requirements

<u>Nutrient</u>	<u>Stage</u>	<u>Amount</u>	<u>Unit</u>
N	Basal	35	kg/ha
P ₂ O ₅	Basal	30	kg/ha
K ₂ O	Basal	35	kg/ha
N	Topdress	25	kg/ha
K ₂ O	Topdress	20	kg/ha

Field images



Varieties

<u>Name</u>	<u>Country</u>	<u>Maturity (days)</u>	<u>Traits</u>
Hollow Crown	KE	120	Long white roots for cool zones.

Fertilizer recommendations

<u>Stage</u>	<u>Product</u>	<u>Rate</u>	<u>Notes</u>
Planting	Well-rotted manure	5000	Improve soil structure before Parsnip planting.
Vegetative growth	CAN	100	Split topdress for Parsnip production.

Pests and diseases

<u>Name</u>	<u>Type</u>	<u>Symptoms</u>	<u>Management</u>
Root maggots	pest	Tunneling and feeding damage on roots.	Rotate crops and maintain field sanitation.
Root cracking and rot	disease	Cracked or rotting roots under uneven moisture.	Maintain even moisture and use well-drained soils.

Yields

<u>System</u>	<u>Typical</u>	<u>Min</u>	<u>Max</u>	<u>Notes</u>
Managed fresh-market production	16	11.2	24	Typical marketable Parsnip yield under irrigated or well-managed conditions.

Season calendars

<u>Country</u>	<u>Region</u>	<u>Planting</u>	<u>Harvest</u>
KE	Highland Vegetable Zones	Mar-Apr or Oct-Nov	Year-round depending on irrigation

Region suitability

<u>Country</u>	<u>Region</u>	<u>Suitability</u>
KE	Highland Vegetable Zones	High

Source: **FarmLens Ltd** - farmlens.africa and app.farmlens.africa. Headquarters: Nairobi, Kenya. This guide was generated from the FarmLens database.