

FarmLens Ltd

Website: farmlens.africa | App: app.farmlens.africa | Headquarters: Nairobi, Kenya



Crop details

Raspberry

Rubus idaeus

Family: Rosaceae

Categories

Fruits & Nuts

Generated: 2026-04-11 06:48

Quick stats

Family	Rosaceae
Typical harvest	6.0 t/ha
Varieties	1
Pests and diseases	2
Seasons	1

Crop profile

Growth habit	shrub
Days to harvest	240
Main uses	Fresh berries, freezing, and premium dessert markets.
Pollination	insect
Origin and where it grows	Suitable for cool highland berry production.

Weather, soil and spacing

Best temperature	10 - 22 °C
Rainfall	800 - 1600 mm/yr
Altitude	0 - 2600 m
Best pH	6 - 7
Soil type	Deep, fertile, well-drained soil with good organic matter.
Row spacing	500 cm
Plant spacing	500 cm
Planting depth	30 cm
Seed rate	kg/ha (check local recommendation)
Nursery days	180

Simple notes for farmers

About the crop: This crop has a growth habit described as "shrub". Harvest typically starts about 240 days after planting.

Main use: Farmers mostly grow this crop for fresh berries, freezing, and premium dessert markets..

Pollination: Mainly insect; healthy flowers and pollinators improve fruit set.

Where it grows: Suitable for cool highland berry production.. Grouped under: Fruits & Nuts.

Best climate: 10 - 22 °C; 800 - 1600 mm/yr; up to about 2600 m a.s.l.

Soil: Best at pH 6 - 7; deep, fertile, well-drained soil with good organic matter..

Farmer guide (Mwongozo wa Mkulima)

<u>Planting</u>	Plant healthy Raspberry seedlings or grafted plants at onset of rains in prepared pits.
<u>Transplanting</u>	Handle Raspberry rootballs carefully and water well after planting.
<u>Irrigation</u>	Maintain steady moisture during establishment and fruit development of Raspberry.
<u>Fertigation</u>	Split nutrients through active growth and fruit development in Raspberry.
<u>Pest scouting</u>	Scout Raspberry regularly for fruit flies, scales, borers, and foliar diseases.
<u>Pruning and training</u>	Prune Raspberry to maintain canopy light, height, and sanitation.
<u>Harvest</u>	Harvest Raspberry at the right maturity stage for fresh-market quality.
<u>Postharvest</u>	Keep Raspberry shaded and cool after harvest to preserve market quality.


Nutrient schedule (Mbolea kwa Hatua)

#	Stage	DAP	Product	Rate	Targets (kg/ha)	Notes
1	Planting or season start	0	Well-rotted manure	8000 kg/ha	N: N/A, P?O?: N/A, K?O: N/A	Improve rooting environment for Raspberry.
2	Pre-fruiting	90	NPK 17-17-17	200 kg/ha	N: 34, P?O?: 34, K?O: 34	Balanced nutrient support for Raspberry.

Nutrient requirements

Nutrient	Stage	Amount	Unit
N	Basal	40	kg/ha
P?O?	Basal	30	kg/ha
K?O	Basal	40	kg/ha
N	Fruiting	40	kg/ha
K?O	Fruiting	50	kg/ha

Field images



Varieties

Name	Country	Maturity (days)	Traits
Heritage	KE	240	Reliable berry canes for cool climates.

Fertilizer recommendations

<u>Stage</u>	<u>Product</u>	<u>Rate</u>	<u>Notes</u>
Planting	Well-rotted manure	8000	Organic matter for Raspberry establishment.
Pre-fruiting	NPK 17-17-17	200	Balanced fertilizer ahead of major Raspberry crop load.

Pests and diseases

<u>Name</u>	<u>Type</u>	<u>Symptoms</u>	<u>Management</u>
Fruit flies	pest	Stings, larval feeding, and rotting fruits.	Field sanitation, trapping, and timely harvest.
Anthraxnose and fruit rots	disease	Lesions on flowers, leaves, or fruits reducing quality.	Prune for airflow, keep orchards clean, and protect during wet periods.

Yields

<u>System</u>	<u>Typical</u>	<u>Min</u>	<u>Max</u>	<u>Notes</u>
Managed orchard production	6	3.6	9.6	Typical orchard yield for Raspberry under practical management.

Season calendars

<u>Country</u>	<u>Region</u>	<u>Planting</u>	<u>Harvest</u>
KE	Managed Orchard Zones	Mar-Apr or Oct-Nov	Depends on variety and agroecology

Region suitability

<u>Country</u>	<u>Region</u>	<u>Suitability</u>
KE	Managed Orchard Zones	Medium

Source: **FarmLens Ltd** - farmlens.africa and app.farmlens.africa. Headquarters: Nairobi, Kenya. This guide was generated from the FarmLens database.