

FarmLens Ltd

Website: farmlens.africa | App: app.farmlens.africa | Headquarters: Nairobi, Kenya



Crop details

Brussels Sprouts

Brassica oleracea var. gemmifera

Family: Brassicaceae

Categories

Vegetables

Generated: 2026-04-11 04:46

Quick stats

Family	Brassicaceae
Typical harvest	9.0 t/ha
Varieties	1
Pests and diseases	2
Seasons	1

Weather, soil and spacing

Best temperature	10 - 20 °C
Rainfall	450 - 800 mm/yr
Altitude	800 - 2800 m
Best pH	6 - 7
Soil type	Fertile well-drained loam rich in organic matter.
Row spacing	40 cm
Plant spacing	20 cm
Planting depth	1.5 cm
Seed rate	4 kg/ha

Crop profile

Growth habit	annual
Days to harvest	120
Main uses	Mini head vegetable for fresh and premium retail chains.
Pollination	insect
Origin and where it grows	Niche cool-season crop in East African highland horticulture.

Simple notes for farmers

About the crop: This crop is annual; it grows and is harvested in one season. Harvest typically starts about 120 days after planting.

Main use: Farmers mostly grow this crop for mini head vegetable for fresh and premium retail chains..

Pollination: Mainly insect; healthy flowers and pollinators improve fruit set.

Where it grows: Niche cool-season crop in East African highland horticulture.. Grouped under: Vegetables.

Best climate: 10 - 20 °C; 450 - 800 mm/yr; up to about 2800 m a.s.l.

Soil: Best at pH 6 - 7; fertile well-drained loam rich in organic matter..

Farmer guide (Mwongozo wa Mkulima)

<u>Planting</u>	Establish Brussels Sprouts in a fine weed-free seedbed and keep emergence moisture steady.
<u>Transplanting</u>	Direct seed or transplant depending on production system.
<u>Irrigation</u>	Maintain even soil moisture for steady Brussels Sprouts growth and quality.
<u>Fertigation</u>	Use split nitrogen and potassium for market-quality Brussels Sprouts.
<u>Pest scouting</u>	Scout Brussels Sprouts weekly for chewing pests, sap suckers, and foliar diseases.
<u>Pruning and training</u>	No pruning required unless sanitation or staking is needed.
<u>Harvest</u>	Harvest Brussels Sprouts at market maturity for the intended use.
<u>Postharvest</u>	Cool and shade Brussels Sprouts promptly after harvest.

Nutrient schedule (Mbolea kwa Hatua)

<u>#</u>	<u>Stage</u>	<u>DAP</u>	<u>Product</u>	<u>Rate</u>	<u>Targets (kg/ha)</u>	<u>Notes</u>
1	Basal	0	NPK 17-17-17	200 kg/ha	N: 34, P ₂ O ₅ : 34, K ₂ O: 34	Basal fertilizer for Brussels Sprouts.
2	Topdress	21	CAN	100 kg/ha	N: 26, P ₂ O ₅ : N/A, K ₂ O: N/A	Support active Brussels Sprouts vegetative growth.

Nutrient requirements

<u>Nutrient</u>	<u>Stage</u>	<u>Amount</u>	<u>Unit</u>
N	Basal	35	kg/ha
P ₂ O ₅	Basal	30	kg/ha
K ₂ O	Basal	35	kg/ha
N	Topdress	25	kg/ha
K ₂ O	Topdress	20	kg/ha

Field images



Varieties

<u>Name</u>	<u>Country</u>	<u>Maturity (days)</u>	<u>Traits</u>
Long Island Improved	KE	120	Compact sprouts on upright stems.

Fertilizer recommendations

<u>Stage</u>	<u>Product</u>	<u>Rate</u>	<u>Notes</u>
Planting	Well-rotted manure	5000	Improve soil structure before Brussels Sprouts planting.
Vegetative growth	CAN	100	Split topdress for Brussels Sprouts production.

Pests and diseases

<u>Name</u>	<u>Type</u>	<u>Symptoms</u>	<u>Management</u>
Aphids	pest	Leaf curling and sticky honeydew.	Scout frequently and use selective control when pressure builds.
Leaf spot complex	disease	Necrotic spotting and reduced leaf quality.	Improve airflow, rotate crops, and avoid prolonged leaf wetness.

Yields

<u>System</u>	<u>Typical</u>	<u>Min</u>	<u>Max</u>	<u>Notes</u>
Managed fresh-market production	9	6.3	13.5	Typical marketable Brussels Sprouts yield under irrigated or well-managed conditions.

Season calendars

<u>Country</u>	<u>Region</u>	<u>Planting</u>	<u>Harvest</u>
KE	Highland Vegetable Zones	Mar-Apr or Oct-Nov	Year-round depending on irrigation

Region suitability

<u>Country</u>	<u>Region</u>	<u>Suitability</u>
KE	Highland Vegetable Zones	High

Source: **FarmLens Ltd** - farmlens.africa and app.farmlens.africa. Headquarters: Nairobi, Kenya. This guide was generated from the FarmLens database.